



Published in final edited form as:

Clin Sports Med. 2011 January ; 30(1): 145–163. doi:10.1016/j.csm.2010.09.006.

Helmets and Mouth Guards: The Role of Personal Equipment in Preventing Sport-Related Concussions

Daniel H. Daneshvar, MA^a, Christine M. Baugh^b, Christopher J. Nowinski^{c,d}, Ann C. McKee^{c,i}, Robert A. Stern, PhD^{c,k}, and Robert C. Cantu, MD^{c,e,f,g,h,i,l}

^aMD / PhD Graduate Student, Boston University School of Medicine (BUSM), Boston, MA

^bResearch Coordinator, Center for the Study of Traumatic Encephalopathy (CSTE), BUSM, Boston, MA

^cCo-Director, CSTE, BUSM, Boston, MA

^dCo-Founder and President, Sports Legacy Institute, Waltham, MA

^eClinical Professor, Department of Neurosurgery, BUSM, Boston, MA

^fChief of Neurosurgery Service, Emerson Hospital, Concord, MA

^gChairman, Department of Surgery, Emerson Hospital, Concord, MA

^hDirector of Sports Medicine, Emerson Hospital, Concord, MA

ⁱCo-Director, Neurologic Sports Injury Center, Brigham and Women's Hospital, Boston, MA

^jAssociate Professor of Neurology and Pathology, BUSM, Boston, MA

^kAssociate Professor of Neurology, BUSM, Boston, MA

^lCo-Founder, Sports Legacy Institute, Waltham, MA

© 2010 Elsevier Inc. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors have nothing to disclose

| | | | | | |
|---------------------------|---------------------------|---------------------------|-----------------------|---------------------------|--|
| Daniel H. Daneshvar, MA | Christine Baugh | Christopher J. Nowinski | Ann C. McKee, MD | Robert A. Stern, PhD | Robert C. Cantu, MD |
| CSTE | CSTE | CSTE | CSTE | CSTE | CSTE and Department of Neurosurgery |
| BUSM | BUSM | BUSM | Bedford VA | BUSM | BUSM |
| 72 East Concord St, B7800 | 72 East Concord St, B7800 | 72 East Concord St, B7800 | 200 Springs Rd, 182-B | 72 East Concord St, B7800 | John Cuming Building, Suite 820, 131 ORNAC |
| Boston, MA 02118 | Boston, MA 02118 | Boston, MA 02118 | Boston, MA 01730 | Boston, MA 02118 | Concord, MA 01742 |
| (617) 414-8382 | (617) 638-6143 | | (781)687-2913 | (617)638-5678 | (978) 369-1386 |
| (617) 414-1197 (fax) | (617) 414-1197 (fax) | (617) 414-1197 (fax) | (781)687-3515 (fax) | (617) 414-1197 (fax) | (978) 287-0047 (fax) |
| ddanesh@bu.edu | cbaugh@bu.edu | nowinski@bu.edu | amckee@bu.edu | bobstern@bu.edu | rcantu@emersonhosp.org |

Abstract

Every year, millions of athletes in the United States experience concussions. With athletes at all levels of play getting bigger, faster, and stronger, it has been suggested that newer technologies may provide an opportunity to reduce the risk and severity of these all too frequent injuries. Although helmets have been shown to decrease the rate of catastrophic head injuries, and mouth guards have decreased the risk of dental and oral injuries, the protective effect of helmets and mouth guards on concussions has not been conclusively demonstrated. In this review, the current literature pertaining to the effect that equipment has on concussions is evaluated. Understanding the role that this equipment plays in preventing concussions is complicated by many factors, such as selection bias in non-randomized studies, variations in playing style, and risk compensation in sports with mandatory protective equipment. At this point, there is little evidence supporting the use of specific helmets or mouth guards to prevent concussions outside of specific sports such as cycling, skiing, and snowboarding. Improving coach and player education about proper concussion management, encouraging neck strengthening exercises, and minimizing high-risk impacts may provide a more fruitful avenue to reduce concussions in sports.

Keywords

concussion; equipment; helmet; headgear; mouth guard; face shield; sport

INTRODUCTION

Because the brain is freely floating within the cerebrospinal fluid, it moves at a different rate than the skull in response to a collision (1). This discrepancy can result in a collision between the brain and skull, either on the side of the impact, coup, or opposite the impact, contrecoup (2). The high speed deceleration associated with these impacts may also result in stretching of the long axons at the base of the brain, resulting in diffuse axonal injury (3). Depending on the extent of these injuries, neurologic dysfunction may be observed (4).

Every year, approximately 1.7 million people in the United States are hospitalized or die as a result of a traumatic brain injury (TBI) (5). These figures, however, are believed to drastically under represent the total incidence of TBI, as many individuals with mild or moderate TBI do not seek medical care (5,6). A portion of these brain injuries are considered concussions, meaning that a direct or indirect blow to the head, face, neck, or body results in an alteration of mental status or produces one or more of 25 recognized post-concussion symptoms (7). It has been estimated that 1.6 million to 3.8 million of these concussions occur annually as a direct result of participation in athletics (5,6,8). The changes in neurologic function associated with concussion often present rapidly and resolve spontaneously (9). As such, many concussions are unreported and unrecognized by coaches, trainers, or the athletes themselves (10–13). A further confounding factor resulting in underreporting the total incidence of concussions is the desire of the athlete to return to play (14).

Most symptoms associated with concussion are transient; however, there are several ways in which concussions can have lasting symptoms. For example, in some cases of concussion, memory impairment has been shown to last for months (4). Furthermore, post-concussion syndrome (PCS) may occur, especially in situations where an athlete is not properly treated following a concussion. PCS presents with physical, cognitive, emotional, and behavioral symptoms that can take months or even years to resolve (15,16). If an athlete returns to play before symptoms resolve, the athlete risks a rare but sometimes fatal event known as second impact syndrome (17,18). Additionally, repetitive concussive and subconcussive blows can

cause Chronic Traumatic Encephalopathy (CTE) or Chronic Traumatic Encephalomyelopathy (CTEM) (19,20).

The importance of understanding and preventing these impacts is increasing, and Athletes have been getting bigger, faster, and stronger, leading to more forceful collisions, which are more likely to cause concussions (21,22). The mechanisms underlying these concussions, as well as methods of prevention, have been investigated both in the lab and in the field. The simplest of these preventative measures appear to be rule changes, rule enforcement, and player and coach education (10). In addition to these suggestions, equipment changes have been proposed in an attempt to help prevent concussions, including modifications of helmets and mouth guards. This equipment has been critical for injury prevention; helmets have been shown to protect against skull fracture, severe traumatic brain injury and death, while mouth guards protect against oral and dental injury (23–25). However, the specific effects of helmets and mouth guards on concussion incidence and concussion severity are less clear.

HELMETS AND HEADGEAR

Protective headgear and helmets decrease the potential for severe TBI following a collision by reducing the acceleration of the head upon impact, thereby decreasing both the brain-skull collision, as well as the sudden deceleration induced axonal injury (26). The energy absorbing material within a helmet accomplishes this by compressing to absorb force during the collision and slowly restoring to its original shape. This compression and restoration has the effect of prolonging the duration of the collision while reducing the total momentum transferred to the head (27). There is quite a bit of variation in helmet design based on the demands and constraints of each sport. While helmets and headgear in most sports are quite good at mediating the high impact collisions responsible for severe TBI, the question remains as to what extent the helmets and headgear of each sport are able to respond to the lower impact collisions responsible for concussion.

American Football

Helmet Design—There has been a great deal of focus on the protection afforded by helmets in football. The primary intent of early football helmets, first reported in use during an Army-Navy game in 1893 and constantly evolving throughout the 1900s, was to prevent catastrophic head injury and the resultant morbidity and mortality (28). These early helmets, then nothing more than leather padding, were slowly phased out as metal and plastics were added to provide additional protection. However, even these basic helmets were not required for college play until 1939, and were not mandated until 1940 for athletes in the National Football League (NFL) (28). Despite these innovations throughout the early twentieth century, the incidence of head injuries continued to increase prompting the formation of the National Operating Committee on Standards for Athletic Equipment (NOCSAE), to initiate research efforts for head protection in 1969 and to implement the first football helmet safety standards in 1973 (29).

These initial NOCSAE guidelines, the framework of which are still in existence today, were meant to develop a standard method for measuring a particular helmet's ability to endure the annual repetitive impacts associated with football in conditions as varied as freezing cold, driving rain, heavy snow, or high heat and humidity (28,30). However, because football collisions at that time were responsible for cerebral hematoma, cervical fractures, and death, the primary concern was the helmet's response to the most acutely severe, linear acceleration inducing impacts, rather than its response to the wide range and types of force that could result in concussion (31–33). While this focus, along with rule changes, ultimately proved successful in decreasing the risk of head injury, the hard-shelled helmet that resulted may not be best suited for protecting against the lower forces that also include a component of rotational

acceleration, which are believed to cause the majority of concussions (23,27,34,35). Furthermore, there is evidence that newly proposed helmet testing methods, meant to encourage the development of helmets better suited to protecting against concussions, may be less accurate than the current testing methods at simulating in vivo concussive impacts (36).

Mechanism of Concussion—Many studies have attempted to measure the forces associated with different types of head impacts. While most early attempts relied on sensors housed within the helmet itself, modern studies have used sensors in contact with the athlete's head (37,38). This new methodology has proven more accurate, as helmet acceleration does not always accurately reflect the acceleration of the head itself (39). These studies have helped develop a better understanding of the types of impacts associated with concussion.

One modern study, examining 19,224 high school football impacts during 55 practices and 13 games, found that impacts to the top of the head were associated with the highest force and the shortest duration of impact, resulting in the highest head jerk (40). Although these impacts to the top of the head are more associated with severe injuries to the cervical spinal cord, rotational acceleration is more closely linked to concussion (41,42). Impacts to the front of the head resulted in the highest rotational acceleration and were the most frequent amongst the high schoolers (40). A similar study following 72 collegiate football players found these impacts to the top of the head 10% less frequent, and associated with 1–2g lower accelerations, than lateral hits, perhaps reflecting improved tackling form or increased neck strength (43). These findings, of fewer hits to the top of the head in collegiate athletes as compared to high school athletes, have been validated elsewhere and reported in studies of professional athletes as well (27, 44). Biomechanical reconstruction of recorded concussive impacts in NFL athletes underscore both the large role that rotational acceleration plays in concussion, as well as the importance of neck strength in mitigating this rotation (45). This method of experimentally replicating recorded collisions may provide additional information about the relationship between both linear and angular head acceleration and injury outcome, resulting in better helmet designs (26,45). However, more work is needed to ensure that these models properly replicate factors potentially influencing concussion like bracing, neck strength, and body tension (46).

Helmet Designs—Football helmets have evolved from little more than relatively modest leather headgear to the modern designs incorporating metals, plastics, and rubber. Early analyses revealed that helmets with pneumatic padding within suspension liners were most effective at absorbing the high intensity impacts that were of early concern (32,33). Modern football helmets incorporate similar features, with hard plastic exteriors housing materials of various stiffness to absorb the force of collision and an inflating system meant to ensure proper fit (28). Football helmets incorporate these basic design elements in various ways, in an attempt to afford better protection to the wearer.

There are several basic designs commonly used in the NFL. One helmet design relies on a continuous tube-like inflatable air fit system, nested into a molded foam network consisting of two different foams with different absorptive properties, ethyl vinyl acetate and polyvinyl chloride nitrile rubber (vinyl nitrile). Front and back pads consist of similar inflatable systems. A second design uses die-cut, rather than molded, foams placed into a case. This creates a laminar system of foam, into which air can be introduced to ensure proper fit. An interchangeable molded urethane front pad completes the system. A third design uses a different approach: although the air-liner is similar to the second system, it is not incorporated into the foam components. Instead, a foam molded ethyl vinyl acetate component with vinyl nitrile inserts, similar to helmet 1, separates the air-liner fit system from an inner shell of expanded polypropylene. Additionally, the plastic outer shell is ventilated and lighter than the previous two. A non-interchangeable vinyl nitrile front pad completes the padding (28).

Recently, helmet manufacturers have begun to design helmets specifically intended to protect against concussion. A description of several of these helmets is in Table 1. One such helmet incorporates distinct design features meant to improve energy attenuation in response to lateral blows. These features include: an exterior shell extending anterior to and distal to traditional shell shapes along the wearer's mandible, increased offset from the interior surface of the shell to the wearer's head in this area, and a unique interior liner construction (47). One newer helmet design creates air turbulence within specialized shock absorbers to allow for differential response to a wider range of impacts levels (48). However, the effect of these and other changes must be further studied, both in the lab and on the field.

Evaluating Helmets—Unfortunately, there have been few studies evaluating the effectiveness of different helmet designs in reducing concussions. The studies that have taken place have tended to be non-randomized, retrospective analyses, and have thus suffered from the same general pitfalls including selection bias and overreliance on subject recollection. Additionally, because the guidelines for concussion assessment were not easily available until recently, many studies relying on coaches and athletic trainers to diagnose concussions may have drastically underreported the total number of concussions as only the most severe injuries would have been counted (49,50). Even after educational outreach efforts by the CDC, the Sports Legacy Institute, and others, knowledge about current concussion guidelines remains an issue (51,52). By only reporting on the relationship between helmets and the most severe concussive injuries, these types of studies risk ignoring the vast majority of concussions, which fall into the mild and moderate range (38). Studies relying on hospitalization data would have a similar bias.

Data collected by the National Athletic Injury/Illness Reporting system from a sample of high school and collegiate athletes during the 1975-77 seasons showed no difference in concussion rate among 13 helmet designs. However, this study was hampered by poor concussion reporting, as a rate of only one concussion per 10,000 athlete exposures was reported, less than one-fourth the most conservative current rates (53,54).

Data on helmet models used and occurrence of cerebral concussions over five seasons were collected from a representative sample of college football teams, consisting of a total of 8,312 player-seasons and 618,596 athlete-exposures. Of the ten models of football helmets included in the analyses, the Riddell M155 had a significantly lower than expected frequency of concussions, whereas the Bike Air Power had a significantly higher frequency (55). However, the number of concussions was again less than half of current conservative estimates of concussion rates (54). Additionally, over half of the athletic exposures in the two statistically significant helmet models were from schools with over 97% of the same helmet type, exposing the study to a potential bias. Because the study relied on athletic trainer report, and all but a few athletes at the majority of schools had the same helmet type, differences in athletic trainer reporting could have had significant effects on the observed concussion rates for many of the athletes studied. To directly compare helmets, each school, and thus each athletic trainer, would have needed to have a mix of all ten of the helmet types studied. The reported relationship is further obscured by the fact that there is a near linear relationship between the number of athletic-exposures and the rate of concussions reported; the helmets with the most exposures tended to have the highest rate of concussion exposure. This finding could be in part due to the fact that larger programs, with more athletes in total and more of the most popular helmet types, may have had staff better trained to recognize concussions.

Another study evaluated the effect of polyurethane helmet covers. Over three seasons from 1992-94, a survey of 155 athletes, identified as having purchased a polyurethane helmet cover in the previous year, were given detailed surveys related to their concussion history in the seasons before and after using the device. Athletes who reported more concussions in the four

years prior to adopting the cover also reported a higher rate of concussion reoccurrence while using the device (56). These results reflect the findings of other studies, which report that players with a history of concussion are significantly more likely to suffer a new concussion as those with no previous history (34,55). Therefore, the use of a polyurethane football helmet cover does not appear to provide additional protection against incurring concussions in the future. In a more recent study, one cohort of 1,173 high school athletes given Riddell Revolution helmets were compared to 968 using standard helmets. All athletes were given a baseline Immediate Post-Concussion Assessment and Cognitive Testing (ImPACT) examination. Over the following three years, whenever an athlete experienced a potential concussive blow, he was assessed for concussive signs using the ImPACT test. During the course of the study, the concussion rate in athletes wearing the Revolution helmet was 5.3% as compared to a concussion rate of 7.6% in athletes wearing the standard helmet [$\chi^2(1, 2, 141) = 4.96, P < 0.027$]. Athletes wearing the Revolution helmet appeared to have a 31% decreased relative risk for sustaining a diagnosed concussion as compared to those who were not (47). Limitations in the study design diminish the strength of the findings, however. The players in the Revolution helmets had new helmets while the standard helmets were of varying age, and helmets tend to become less effective over time. In addition, because athletic trainers are often unaware of all but the most severe concussions, the results may have been influenced more by reporting rates of the high school staff than absolute differences in risk of concussion. Furthermore, concussions were diagnosed by ImPACT instead of a combination of neurological exam that includes balance testing. Finally, the study was funded by Riddell, and a Riddell employee was a lead author which poses an obvious conflict of interest. Although there was a difference in rate of concussion between the two groups, concussed athletes in the two groups did not differ significantly in their average number of days to recover and return to play following their concussions.

Baseball

Several new “concussion-proof” helmet designs have been proposed and are now being introduced into Major and Minor League Baseball, but their degree of effectiveness in preventing concussion has not yet been demonstrated in any epidemiologic study. However, the role of standard baseball helmets in preventing more serious head injuries has been well validated (57).

Cycling

Helmets have been long shown to decrease the rates of head injury in cyclists (58–61). However, there have been no studies evaluating the different bicycle helmet designs in response to concussive impacts.

Ice Hockey

Ice Hockey, like football, is a helmeted sport associated with a significant risk of concussion. For aforementioned reasons, including increased awareness of diagnostic criteria, the rate of diagnosed concussions has been increasing; nonetheless, helmets are largely responsible for protecting hockey players from the most catastrophic head injuries (10,11,54). The introduction of ice hockey helmet standards and proper utilization of helmets have resulted in a decrease in fatal and catastrophic head injuries, but with an increase in rate of concussions (62). While this increase in rate is likely due in large part to better concussion awareness and recognition, aggressive play is likely also responsible for this increase in rate (63).

Concussions in hockey most commonly occur as a result of collision with an opponent or collision with the boards (64). Measurements from sensors within the helmets of hockey players have found that the impacts sustained by hockey players are comparable in magnitude to those experienced by football linemen, but occur at approximately one-third of the frequency (38).

Since helmets have been made mandatory in hockey, there has been little published comparing the protective effects of different hockey helmets. However, there is evidence that player flexion and anticipation prior to a collision decreases the risk of concussion, providing a potential avenue for future helmet design (46). Additionally, newer helmet testing methods have now begun to take into account the rotational acceleration component involved in a collision, which better simulates concussive impacts (35).

Lacrosse

In a recent study of athletic trainers, concussion was found to be the most common injury in lacrosse, and was responsible for a higher percentage of all injuries among boys (73%) and men (85%) than among girls (40%) and women (41%). In men, the primary injury mechanism was player-to-player contact, whereas women's injuries primarily resulted from stick or ball contact (65).

Although the rate of concussions has increased dramatically in many sports, some have argued that this observation in men's lacrosse may be, in part, explained by the introduction of a new helmet. One study compared the rate of concussion in the years immediately following the helmet's introduction (1996–97 to 2003–04) to the preceding years (1988–89 to 1995–96). In practices, the rate of concussion increased by 0.14 concussions per 1000 A–E (95% CI = 0.09, 0.19, $P < .01$). In games, the rate increased by 0.84 (95% CI = 0.52, 1.16, $P < .01$) (66). However, this increase is certainly due, in part, to improved detection and diagnosis of concussion during that time frame.

Rugby

Headgear in rugby consists of relatively sparse padding, and its use is not mandated. As a result, the role of headgear in preventing concussion and head injuries can be more easily studied. In one prospective study of 294 players under age 15, headgear was distributed to players. Over the course of the study, there were 1179 player exposures with headgear and 357 without headgear. During this period, there were only nine reported concussions, seven of which occurred to players wearing headgear, two to those not wearing headgear. As a result, there was no evidence indicating a protective effect of headgear; in fact these data showed headgear having a non significant deleterious effect (67). However, as only concussions that were medically verified were reported, many minor concussions may have been unreported. Additionally, headgear use was not randomized, as athletes had the choice of whether or not to wear headgear. This produces a potential bias, as athletes more concerned about head injury are more likely to wear headgear; certainly a subset of these athletes had had prior concussions and was therefore more susceptible to having additional concussions. In another study of 304 rugby players, followed weekly, headgear was shown to have had a non significant protective effect on concussions, but a significant protective effect on orofacial and scalp injuries (68). However, only 22 concussions were recorded and the study was not adequately powered to determine a suitable effect size. Additionally, one survey-based study of 131 men's club rugby union participants from eight university teams in the United States reported 76 total concussed athletes. Although 51% of all of the surveyed athletes were not wearing headgear, 76% of all concussed athletes reported not wearing headgear. The remaining who were concussed while wearing headgear reported that their concussions were less severe, as compared to the reports of those who were not wearing headgear (69).

However, in the most thorough study, 1493 participants from four rugby leagues (under 13, under 15, under 18 and under 20) were randomly assigned one of two types of headgear, or no headgear, and followed over two years. Although compliance to the random assignment was relatively low, nearly half of all athlete-exposures consisted of athletes wearing one of the two

headgear types. Regardless, the use of these padded headgears did not affect the rate of concussion, nor the number of days missed as a result of concussion (70).

These findings, which indicate that rugby headgear does not appear to have a protective effect in concussion prevention, correspond to laboratory findings indicating that headgear are maximally compressed at impacts far below those likely to cause a concussion (71). Because they are unable to absorb additional force well below the threshold at which concussions occur, they would not be expected to have a major effect on the incidence of concussion. However, this ceiling effect may be avoided in future headgear by methods such as modifying padding materials and increasing padding thickness (72).

Besides the apparent lack of scientific justification supporting the use of rugby headgear for concussion prevention, additional barriers remain to widespread headgear adoption. Although athletes tend to report that headgear is beneficial, athletes in one survey commonly reported that wearing headgear was uncomfortable, poorly ventilated, and often grabbed by opponents during play (69,73).

Soccer

Several studies have sought to measure the nature of impacts experienced by soccer players. It was found that, although soccer players experienced significantly fewer impacts per hour than high school football lineman or hockey players, each impact tended to be associated with higher accelerations; whereas 20% of impacts in soccer were over 75g, approximately 5% of impacts were over 75g in football lineman and hockey players (38). However, the head-ball collisions that were studied are not the most common source of concussions in soccer. The majority of concussions in soccer occur during the act of heading as a result of head-head or head-arm collisions (74). Another study following athletes over three years found that the most common site of impact in a concussion causing collision was the temporal area of the head (44). This finding supports those of other sports and underscores the importance of rotational acceleration in concussion.

There are various types of headgear proposed to limit the effect of concussion in soccer athletes. While there have been non-randomized studies of the effect of headgear on head injuries in soccer, there have been few analytic studies looking specifically at the role of headgear on concussion rate or severity (44,75). This remains a potential area of interest.

One retrospective anonymous online survey of youth soccer players, aged 12 to 17 years old, studied the role of headgear on concussion symptoms. Of those eligible for the study, 216 athletes were included in the non-headgear-wearing group and 52 were included in the headgear-wearing group. This study was strengthened by the fact that it asked respondents not only how many concussions they had experienced in the prior season, but also asked them how many times they had experienced specific symptoms associated with concussions, in response to a collision. Interestingly, 7.2% of all athletes reporting having experienced at least one concussion, whereas 47.8% reported having experienced concussive symptoms at least once. As expected, it was found that athletes wearing headgear were significantly less likely to receive laceration to those areas of the scalp and face covered by the headgear. Interestingly, although the group that chose to wear headgear reported having experienced more concussions prior to the study (42.3% had experienced at least one prior concussion and 26.9% had experienced more than one, compared to 11.1% and 4.6% respectively) and would therefore be expected to more knowledgeable and at increased risk of having an additional concussion, it was found that not wearing headgear was associated with a 2.65 relative risk of concussion ($p < 0.0001$) (75). This finding might be explained in part by the fact that fewer athletes who wore headgear considered themselves to be “a header” (44.2% of those who wore headgear as compared to 51.4% of those who did not wear headgear). Although this study is quite

promising, it was not ideal because headgear use was non-randomized, and the retrospective study relied principally on information recollected by athletes at the end of season. Although the anonymous nature of the survey meant that it was not possible to ask follow up questions of participants or verify data, this is nonetheless a likely strength as many studies have found more accurate concussion data when athletes anonymously report their history. However, if all soccer players were to wear headgear, the effect on concussions may be complicated by risk compensation. Rule changes in football, hockey, and lacrosse have suggested that mandating headgear removes inhibitions to strike, or risk strikes to, the head, as it reduces pain from scalp injuries and lacerations. If all players were to become accustomed to a playing style in which contact to the head was no longer 'off limits', the addition of headgear might result in an increase in the frequency of total collisions to the head, and potentially increase the total number of concussions as well.

There is some evidence that the same headgear may not be appropriate for all athletes. One study evaluating heading kinematics found that women experienced higher acceleration than men when heading a ball (76). Although the majority of concussions do not occur as a result of head-ball interactions, these findings nonetheless indicate that gender differences may need to be accounted for in future headgear designs. Additional laboratory testing found that, although headgear is unlikely to be effective in attenuating impact during head-ball collisions, it may decrease the impact of head-head collisions by nearly 33% (77).

However, because there is a specific mechanism of injury associated with concussion, the simplest preventative strategy may not be newer equipment. As concussions typically occurred in both men and women as a result of contact while heading the ball, limiting this type of contact through rule changes may be appropriate (4).

Skiing/Snowboarding

Observational studies have found that 12.1% of US skiers wear helmets (78). However, at the Saint Anthony Central Hospital (Denver, CO) Level I trauma center from 1982 through 1998, only 3 of the total 1,214 patients admitted for all ski-related head injuries were wearing a helmet (79). Several studies argue that helmets may reduce risk of concussion by up to 60% (80–83).

MOUTH GUARDS

During the 1960s and 1970s, the use of mouth guards was made mandatory in many sports including football, ice hockey, lacrosse, field hockey, and boxing. The rationale for these rule changes was to provide additional protection against dental injury, orofacial injury and to reduce a player's risk of concussion (25,84). However, at that time, as well as now, there is little evidence that mouth guards provide protection against concussion.

American Football

There is interest in the possibility that better designed mouth guards might help dissipate force, thereby reducing the magnitude of the impact. Because mouth guards are already mandatory equipment in football, there is an opportunity to evaluate the role of specific types of mouth guards in preventing concussion. There is some scientific rationale that custom fitted mouth guards might be more effective at measurably absorbing the force of impact (85). A study of 28 high school and college football players suggested a decreased rate of concussion following the use of customized mandibular orthotics; however, this study was marked by a number of design flaws. Concussion rate prior to customized mandibular orthotics was measured by self report, whereas concussion rate following orthotic use was only calculated based on concussions diagnosed by athletic trainers and coaches. Also, because all athletes were given orthotics, the observed decrease in concussion rate could simply be an artifact of different styles

or age of play; all athletes were necessarily older when using the custom orthotics than they were when using standard mouth guards. Finally, the rate of concussion prior to custom orthotic use was not limited to games, whereas only concussions during games were counted towards the rate of concussions following custom orthotics (86). No large study has been able to demonstrate a significant difference in type of mouth guard and concussion rate. One study recruited 87 of a total 114 Division 1 teams to participate in a study evaluating the effect of various mouth guard types on rate of concussion. There was no statistically significant result between the different mouth guards (87). These findings have since been replicated by other large, multi-center cohort studies (88).

Basketball

Many are interested in the potential role of mouth guards in reducing the number of concussions in basketball. However, recent studies have found no significant differences in concussion rates between mouth guard users and nonusers (89).

Ice Hockey

There has been some interest in the degree to which helmets, mouth guards, and visors affect concussions in athletes. There are several different hockey helmet designs that have been shown to decrease total impact experimentally; however, there have been no studies to determine their utility in reducing concussions.

Mouth guards are not mandatory in the National Hockey League (NHL) as there is debate over the extent to which they reduce concussion incidence and severity. In a study of 1,033 NHL athletes, the rate of concussion was 1.42 times greater in individuals who did not wear mouth guards, as compared to those who did. However, this difference was not statistically significant (95% CI 0.90 to 2.25). Despite the non-significant finding regarding concussion rate, symptom severity was significantly decreased by use of mouth guards. Symptom severity, measured using the modified McGill ACE symptom scale, was found to be significantly worse in athletes who were not wearing mouth guards than in those who were ($p < 0.01$) (90,91).

Rugby

In rugby, although there is evidence that mouth guards protect against orofacial injuries, a study of 304 rugby players followed weekly showed no significant effect of mouth guards on the incidence of concussions, although only 22 total concussions were observed (68).

Soccer

One anonymous survey of 278 youth soccer players aged 12 to 17 years old found no significant relationship between mouth guard use and the rate of concussion (75).

OTHER EQUIPMENT

Facial Protection

During the 1970s, full facial protection was mandated by all organized youth ice hockey associations worldwide. However, a number of studies have shown no significant relationship between use of visors and concussion rate in high school, college, or NHL athletes (91–94).

In junior A ice hockey, full faceguards were found to provide a 4.7 times reduction in eye injuries and a non-significant reduction in the rate of concussion from 12.2 to 2.9 concussions per 1000 player hours compared with no faceguard (94). However, this study was hampered by restricted data collection for playing time and injuries, which were only recorded during home games. Therefore, injuries from away games were not included in the analysis. Also,

players under the age of 18 years had to wear mandatory full facial protection, whereas players above the age of 18 years could choose to wear a full shield, half shield, or no shield (95). Playing style might have influenced this decision, as players with riskier playing styles may have chosen to play without visors.

In university level ice hockey, the use of full faceguards was found to reduce the number of games missed because of concussion, but not the incidence of concussion, compared with the use of half faceguard (93). Additionally, while the use of full face shields significantly reduced players' risk of sustaining a dental injury, there was no difference in the incidence of concussions between players wearing different shield types. However, players who sustained concussions while wearing half shields required significantly more time before returning to competition than players who sustained concussions wearing full face shields (96). One potential explanation for this finding has to do with helmet placement. Players wearing half shields may have been tilting their helmets back to allow for an unobstructed view below the visor, thereby getting a clearer view but resulting in improper helmet placement. Alternatively, the minor visual impairment from the half shield could have led to an increased number of hits that were not foreseen by the athlete and therefore may have led to a greater number of concussions. Improper helmet alignment would cause less padding on the forehead and a loose chin strap decreasing the protective effect of the helmet. Neither of the aforementioned studies compared the incidence of concussion in athletes with visors to those without. In professional hockey, the use of a visor did not significantly reduce the prevalence of concussions. Visor use did, however, lower the prevalence of eye and non-concussion head injuries (92).

There is some additional interest in the potential of facial protection to mitigate concussion in other sports as well. Interestingly, in biomechanical reconstructions of professional football concussive injuries, impacts to the facemask resulted in the higher rotational accelerations, likely due to the fact that facemasks sit outside the helmet shell and thus have an increased radius of rotation from the base of the head (45). This may help influence future helmet designs.

In baseball, some have proposed that faceguards may reduce the risk of concussion, although this has not yet been studied. In youth baseball, facemasks have been shown to reduce the incidence of oculofacial injury in a non-randomized prospective cohort study (97). Overall, faceguards are also associated with a reduced risk of facial injury (98).

Baseball Balls

In baseball, softer balls have been shown to reduce the risk of head injury compared with standard balls, but this equipment has yet to be studied in light of concussions. Theoretical biomechanical studies have indicated that baseballs with lower mass and less stiffness have a reduced potential for injury (99). Additionally, laboratory studies have shown that reduced-impact balls are less likely to result in head injury and skull fracture (100). In practice, softer baseballs have been found to yield a 28% reduction in the risk of injury (24,98,101).

Playing Surfaces

As the speed of athletes increases, the momentum transfer and impact associated with their collisions increase as well. The surface on which athletes play affects the player's speed and therefore may influence the rate of concussions. In general, synthetic surfaces are harder and result in faster speeds than natural ones (34). Some studies have found that athletes playing on synthetic fields have a higher risk of injury (102). Laboratory studies have found that different fields have different impact attenuation properties, which would certainly be expected to at least influence head-ground collisions (103).

DISCUSSION

Several studies have provided biomechanical evidence of the reduction of impact forces to the brain due to the use of specific headgear or helmets. However, in the majority of sports, these results have not translated into observed differences in rate or severity of concussion. For some sports in which contact with hard surfaces is possible, such as skiing, snowboarding, and cycling, there is evidence that helmets greatly reduce the incidence of head injuries in general; therefore, helmets are an important part of injury prevention and should be recommended in these sports (61,104–106).

One complicating factor associated with helmet and headgear use is that of risk compensation. In many cases, protective equipment can lead to a false sense of security, resulting in a more dangerous style of play (107). Often, the helmet itself may be used to initiate contact. This tendency to promote a more reckless style of play may help explain the higher rate of injury in children and adolescents, as compared to adults (108).

In general, helmets work best when properly utilized. This means that the helmets must be sized appropriately and worn with all straps correctly fastened and all padding in the proper positioning. Problems arise when helmets are older, incorrectly sized, worn improperly, or when padding is underinflated. Inclement weather also has been shown to affect a helmet's ability to absorb impacts. (28). Additionally, neck strength may be important in minimizing risk of concussion in response to an impact. Therefore, rule changes mandating neck strength training, education about proper tackling form including prohibiting spearing, and monitoring player fatigue is warranted (45). Encouraging practices with limited contact may also be an appropriate way to limit concussive blows in football; however, some studies indicate that helmet only practices are associated with similar impacts as those experienced in full contact play (43).

Although mouth guards have been shown to be effective in preventing dental and oro-facial injury, there is currently no evidence that standard or fitted mouth guards decrease the rate or severity of concussions in athletes (85). The bulk of the evidence indicating a potential protective effect of mouth guards on concussion incidence has been based on a limited case series studies and retrospective, non-randomized cross-sectional surveys (91). There is also evidence that mouth guard use does not result in any difference in neurocognitive test performance following concussion (109). In sports such as hockey, there is no evidence that visors play a protective role in preventing or mitigating concussions (91–94).

Many of the studies on the protective effect of equipment on concussive risk have been complicated by retrospective, non-randomized study designs. Individuals may choose to wear specialized protective equipment based on previous injury history, which has been shown to increase risk of future injuries, or because of a risky playing style. The preponderance of evidence appears to indicate that helmets and mouth guards provide a significant benefit in protecting against many catastrophic head, neck, and orofacial injuries. However, there is not yet significant evidence to advocate their effectiveness in preventing concussion. Nonetheless, additional research is needed, both in the laboratory to improve equipment design, and on the field to verify findings epidemiologically. Although newer equipment remains a promising potential tool in minimizing concussion severity and incidence, other methods such as rule changes, improved concussion education, and proper coaching and training may prove more effective in the immediate future.

Acknowledgments

This work was supported by the Boston University Alzheimer's Disease Center NIA P30 AG13846, supplement 0572063345-5, the National Operating Committee on Standards for Athletic Equipment, the National Collegiate

Athletic Association, the National Federation of State High School Associations, the American Football Coaches Association, and the Sports Legacy Institute

References

1. Viano DC, Casson IR, Pellman EJ, Zhang L, King AI, Yang KH. Concussion in professional football: brain responses by finite element analysis: part 9. *Neurosurgery* 2005 Nov;57(5):891–916. discussion 891–916. [PubMed: 16284560]
2. Drew LB, Drew WE. The contrecoup-coup phenomenon: a new understanding of the mechanism of closed head injury. *Neurocrit Care* 2004;1(3):385–390. [PubMed: 16174940]
3. Meythaler JM, Peduzzi JD, Eleftheriou E, Novack TA. Current concepts: diffuse axonal injury-associated traumatic brain injury. *Arch Phys Med Rehabil* 2001 Oct;82(10):1461–1471. [PubMed: 11588754]
4. McCrory, P.; Meeuwisse, W.; Johnston, K., et al. Consensus statement on concussion in sport--the 3rd International Conference on concussion in sport, held in Zurich, November 2008; *J Clin Neurosci*; 2009 Jun. p. 755-763.
5. Faul, M.; Xu, L.; Wald, MM.; Coronado, VG. Traumatic Brain Injury in the United States: Emergency Department Visits, Hospitalizations and Deaths 2002–2006. US Department of Health and Human Services; 2010 Mar.
6. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil* 2006 Sep–Oct;21(5):375–378. [PubMed: 16983222]
7. Concussion (mild traumatic brain injury) and the team physician: a consensus statement. *Med Sci Sports Exerc* 2006 Feb;38(2):395–399. [PubMed: 16531912]
8. Finkelstein, E.; Corso, P.; Miller, T. The Incidence and Economic Burden of Injuries in the United States. New York, NY: Oxford University Press; 2006.
9. Parker TM, Osternig LR, van Donkelaar P, Chou LS. Recovery of cognitive and dynamic motor function following concussion. *Br J Sports Med* 2007 Dec;41(12):868–873. discussion 873. [PubMed: 17517857]
10. Cantu RC, Mueller FO. The prevention of catastrophic head and spine injuries in high school and college sports. *Br J Sports Med* 2009 Dec;43(13):981–986. [PubMed: 19945980]
11. Mueller, FO.; Cantu, RC. National Center for Catastrophic Injury Research; 2008 Spring. Catastrophic Sport Injury Research 26th Annual Report: Fall 1982-Spring 2008.
12. Delaney JS, Lacroix VJ, Leclerc S, Johnston KM. Concussions among university football and soccer players. *Clin J Sport Med* 2002 Nov;12(6):331–338. [PubMed: 12466687]
13. Field M, Collins MW, Lovell MR, Maroon J. Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. *J Pediatr* 2003 May;142(5):546–553. [PubMed: 12756388]
14. Yard EE, Comstock RD. Compliance with return to play guidelines following concussion in US high school athletes, 2005–2008. *Brain Inj* 2009 Oct;23(11):888–898. [PubMed: 20100125]
15. Barlow KM, Crawford S, Stevenson A, Sandhu SS, Belanger F, Dewey D. Epidemiology of postconcussion syndrome in pediatric mild traumatic brain injury. *Pediatrics* 2010 Aug;126(2):e374–e381. [PubMed: 20660554]
16. Cantu RC, Register-Mihalik J, Guskiewicz KW. A Retrospective Analysis of 215 Athletic Moderate to Severe Concussions. *J Physical Medicine and Rehabilitation*. 2010 ((in press)).
17. Cantu R, Gean A. Second impact syndrome in a small SDH: an uncommon catastrophic result of repetitive head injury with a characteristic imaging appearance. *J Neurotrauma*. 2010 Jun 10;
18. Cantu RC. Second-impact syndrome. *Clin Sports Med* 1998 Jan;17(1):37–44. [PubMed: 9475969]
19. McKee AC, Gavett BE, Stern RA, et al. TDP-43 Proteinopathy and Motor Neuron Disease in Chronic Traumatic Encephalopathy. *J Neuropathol Exp Neurol*. 2010 Aug 11;
20. McKee AC, Cantu RC, Nowinski CJ, et al. Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury. *J Neuropathol Exp Neurol* 2009 Jul;68(7):709–735. [PubMed: 19535999]
21. Wennberg RA, Tator CH. National Hockey League reported concussions, 1986-87 to 2001-02. *Can J Neurol Sci* 2003 Aug;30(3):206–209. [PubMed: 12945942]

22. Barth JT, Freeman JR, Broshek DK, Varney RN. Acceleration-Deceleration Sport-Related Concussion: The Gravity of It All. *J Athl Train* 2001 Sep;36(3):253–256. [PubMed: 12937493]
23. Cantu RC, Mueller FO. Brain injury-related fatalities in American football, 1945–1999. *Neurosurgery* 2003 Apr;52(4):846–852. discussion 852–843. [PubMed: 12657180]
24. McIntosh AS, McCrory P. Preventing head and neck injury. *Br J Sports Med* 2005 Jun;39(6):314–318. [PubMed: 15911597]
25. Using mouthguards to reduce the incidence and severity of sports-related oral injuries. *J Am Dent Assoc* 2006 Dec;137(12):1712–1720. quiz 1731. [PubMed: 17138717]
26. Newman JA, Beusenber MC, Shewchenko N, Withnall C, Fournier E. Verification of biomechanical methods employed in a comprehensive study of mild traumatic brain injury and the effectiveness of American football helmets. *J Biomech* 2005 Jul;38(7):1469–1481. [PubMed: 15922758]
27. Pellman EJ, Viano DC, Withnall C, Shewchenko N, Bir CA, Halstead PD. Concussion in professional football: helmet testing to assess impact performance--part 11. *Neurosurgery* 2006 Jan;58(1):78–96. discussion 78–96. [PubMed: 16385332]
28. Levy ML, Ozgur BM, Berry C, Aryan HE, Apuzzo ML. Birth and evolution of the football helmet. *Neurosurgery* 2004 Sep;55(3):656–661. discussion 661–652. [PubMed: 15335433]
29. Bennett, T., editor. *The NFL's Official Encyclopedic History of Professional Football*. 2 ed.. New York: Macmillan; 1977.
30. (NOCSAE) NOCoSfAE. Standard Drop Test Method and Equipment Used in Evaluating the Performance Characteristics of Protective Headgear. NOCSAE; 2004.
31. Robey JM, Blyth CS, Mueller FO. Athletic injuries. Application of epidemiologic methods. *JAMA* 1971 Jul 12;217(2):184–189. [PubMed: 5108781]
32. Bishop PJ, Norman RW, Kozey JW. An evaluation of football helmets under impact conditions. *Am J Sports Med* 1984 May–Jun;12(3):233–236. [PubMed: 6742308]
33. Myers, TJ.; Yoganandan, N.; Sances, A., Jr; Pintar, FA.; Reinartz, J.; Battocletti, JH. *Biomed Mater Eng*. Vol. 3. 1993 Spring. Energy absorption characteristics of football helmets under low and high rates of loading; p. 15-24.
34. Levy ML, Ozgur BM, Berry C, Aryan HE, Apuzzo ML. Analysis and evolution of head injury in football. *Neurosurgery* 2004 Sep;55(3):649–655. [PubMed: 15335432]
35. Kis M, Saunders F, ten Hove MW, Leslie JR. Rotational acceleration measurements--evaluating helmet protection. *Can J Neurol Sci* 2004 Nov;31(4):499–503. [PubMed: 15595255]
36. Gwin JT, Chu JJ, Diamond SG, Halstead PD, Crisco JJ, Greenwald RM. An investigation of the NOCSAE linear impactor test method based on in vivo measures of head impact acceleration in American football. *J Biomech Eng* 2010 Jan;132(1) 011006.
37. Reid SE, Tarkington JA, Epstein HM, O'Dea TJ. Brain tolerance to impact in football. *Surg Gynecol Obstet* 1971 Dec;133(6):929–936. [PubMed: 5117389]
38. Naunheim RS, Standeven J, Richter C, Lewis LM. Comparison of impact data in hockey, football, and soccer. *J Trauma* 2000 May;48(5):938–941. [PubMed: 10823540]
39. Manoogian S, McNeely D, Duma S, Brolinson G, Greenwald R. Head acceleration is less than 10 percent of helmet acceleration in football impacts. *Biomed Sci Instrum* 2006;42:383–388. [PubMed: 16817638]
40. Broglio SP, Sosnoff JJ, Shin S, He X, Alcaraz C, Zimmerman J. Head impacts during high school football: a biomechanical assessment. *J Athl Train* 2009 Jul–Aug;44(4):342–349. [PubMed: 19593415]
41. Rihn JA, Anderson DT, Lamb K, et al. Cervical spine injuries in American football. *Sports Med* 2009;39(9):697–708. [PubMed: 19691361]
42. Broglio SP, Schnebel B, Sosnoff JJ, et al. The Biomechanical Properties of Concussions in High School Football. *Med Sci Sports Exerc*. 2010 Mar 25;
43. Mihalik JP, Bell DR, Marshall SW, Guskiewicz KM. Measurement of head impacts in collegiate football players: an investigation of positional and event-type differences. *Neurosurgery* 2007 Dec; 61(6):1229–1235. discussion 1235. [PubMed: 18162902]
44. Scott Delaney J, Puni V, Rouah F. Mechanisms of injury for concussions in university football, ice hockey, and soccer: a pilot study. *Clin J Sport Med* 2006 Mar;16(2):162–165. [PubMed: 16603887]

45. Viano DC, Casson IR, Pellman EJ. Concussion in professional football: biomechanics of the struck player--part 14. *Neurosurgery* 2007 Aug;61(2):313–327. discussion 327–318. [PubMed: 17762744]
46. Mihalik JP, Blackburn JT, Greenwald RM, Cantu RC, Marshall SW, Guskiewicz KM. Collision type and player anticipation affect head impact severity among youth ice hockey players. *Pediatrics* 2010 Jun;125(6):e1394–e1401. [PubMed: 20478933]
47. Collins M, Lovell MR, Iverson GL, Ide T, Maroon J. Examining concussion rates and return to play in high school football players wearing newer helmet technology: a three-year prospective cohort study. *Neurosurgery* 2006 Feb;58(2):275–286. discussion 275–286. [PubMed: 16462481]
48. Xenith. Xenith Innovation. 2010. <http://www.xenith.com/football/innovation/>
49. Covassin T, Elbin R 3rd, Stiller-Ostrowski JL. Current sport-related concussion teaching and clinical practices of sports medicine professionals. *J Athl Train* 2009 Jul–Aug;44(4):400–404. [PubMed: 19593422]
50. Notebaert AJ, Guskiewicz KM. Current trends in athletic training practice for concussion assessment and management. *J Athl Train* 2005 Oct–Dec;40(4):320–325. [PubMed: 16404454]
51. Sarmiento K, Mitchko J, Klein C, Wong S. Evaluation of the Centers for Disease Control and Prevention's concussion initiative for high school coaches: "Heads Up: Concussion in High School Sports". *J Sch Health* 2010 Mar;80(3):112–118. [PubMed: 20236412]
52. Sawyer RJ, Hamdallah M, White D, Pruzan M, Mitchko J, Huitric M. High school coaches' assessments, intentions to use, and use of a concussion prevention toolkit: Centers for Disease Control and Prevention's heads up: concussion in high school sports. *Health Promot Pract* 2010 Jan;11(1):34–43. [PubMed: 18400880]
53. Clarke KS, Powell JW. Football helmets and neurotrauma--an epidemiological overview of three seasons. *Med Sci Sports* 1979 Summer;11(2):138–145. [PubMed: 491870]
54. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train* 2007 Apr–Jun;42(2):311–319. [PubMed: 17710181]
55. Zemper ED. Analysis of cerebral concussion frequency with the most commonly used models of football helmets. *J Athl Train* 1994 Mar;29(1):44–50. [PubMed: 16558258]
56. Torg JS, Harris SM, Rogers K, Stilwell GJ. Retrospective report on the effectiveness of a polyurethane football helmet cover on the repeated occurrence of cerebral concussions. *Am J Orthop (Belle Mead NJ)* 1999 Feb;28(2):128–132. [PubMed: 10067717]
57. Nicholls RL, Elliott BC, Miller K. Impact injuries in baseball : prevalence, aetiology and the role of equipment performance. *Sports Med* 2004;34(1):17–25. [PubMed: 14715037]
58. Williams M. The protective performance of bicyclists' helmets in accidents. *Accid Anal Prev* 1991 Apr–Jun;23(2–3):119–131. [PubMed: 2029314]
59. Thompson DC, Rivara FP, Thompson R. Helmets for preventing head and facial injuries in bicyclists. *Cochrane Database Syst Rev*. 2000;(2) CD001855.
60. Haileyesus T, Annett JL, Dellinger AM. Cyclists injured while sharing the road with motor vehicles. *Inj Prev* 2007 Jun;13(3):202–206. [PubMed: 17567979]
61. Scheiman S, Moghaddas HS, Bjornstig U, Bylund PO, Saveman BI. Bicycle injury events among older adults in Northern Sweden: a 10-year population based study. *Accid Anal Prev* 2010 Mar;42(2):758–763. [PubMed: 20159104]
62. Biasca N, Wirth S, Tegner Y. The avoidability of head and neck injuries in ice hockey: an historical review. *Br J Sports Med* 2002 Dec;36:410–427. [PubMed: 12453835]
63. Daly PJ, Sim FH, Simonet WT. Ice hockey injuries. A review. *Sports Med* 1990 Aug;10(2):122–131. [PubMed: 2204098]
64. Flik K, Lyman S, Marx RG. American collegiate men's ice hockey: an analysis of injuries. *Am J Sports Med* 2005 Feb;33(2):183–187. [PubMed: 15701603]
65. Lincoln AE, Hinton RY, Almquist JL, Lager SL, Dick RW. Head, face, and eye injuries in scholastic and collegiate lacrosse: a 4-year prospective study. *Am J Sports Med* 2007 Feb;35(2):207–215. [PubMed: 17204586]
66. Dick R, Romani WA, Agel J, Case JG, Marshall SW. Descriptive epidemiology of collegiate men's lacrosse injuries: National Collegiate Athletic Association Injury Surveillance System, 1988–1989 through 2003–2004. *J Athl Train* 2007 Apr–Jun;42(2):255–261. [PubMed: 17710174]

67. McIntosh AS, McCrory P. Effectiveness of headgear in a pilot study of under 15 rugby union football. *Br J Sports Med* 2001 Jun;35(3):167–169. [PubMed: 11375874]
68. Marshall SW, Loomis DP, Waller AE, et al. Evaluation of protective equipment for prevention of injuries in rugby union. *Int J Epidemiol* 2005 Feb;34(1):113–118. [PubMed: 15561749]
69. Kahanov L, Dusa MJ, Wilkinson S, Roberts J. Self-reported headgear use and concussions among collegiate men's rugby union players. *Res Sports Med* 2005 Apr–Jun;13(2):77–89. [PubMed: 16392441]
70. McIntosh AS, McCrory P, Finch CF, Best JP, Chalmers DJ, Wolfe R. Does padded headgear prevent head injury in rugby union football? *Med Sci Sports Exerc* 2009 Feb;41(2):306–313. [PubMed: 19127196]
71. McIntosh AS, McCrory P. Impact energy attenuation performance of football headgear. *Br J Sports Med* 2000 Oct;34(5):337–341. [PubMed: 11049142]
72. McIntosh A, McCrory P, Finch CF. Performance enhanced headgear: a scientific approach to the development of protective headgear. *Br J Sports Med* 2004 Feb;38(1):46–49. [PubMed: 14751945]
73. Pettersen JA. Does rugby headgear prevent concussion? Attitudes of Canadian players and coaches. *Br J Sports Med* 2002 Feb;36(1):19–22. [PubMed: 11867487]
74. Gessel LM, Fields SK, Collins CL, Dick RW, Comstock RD. Concussions among United States high school and collegiate athletes. *J Athl Train* 2007 Oct–Dec;42(4):495–503. [PubMed: 18174937]
75. Delaney JS, Al-Kashmiri A, Drummond R, Correa JA. The effect of protective headgear on head injuries and concussions in adolescent football (soccer) players. *Br J Sports Med* 2008 Feb;42(2):110–115. discussion 115. [PubMed: 17615173]
76. Tierney RT, Higgins M, Caswell SV, et al. Sex differences in head acceleration during heading while wearing soccer headgear. *J Athl Train* 2008 Oct–Dec;43(6):578–584. [PubMed: 19030135]
77. Withnall C, Shewchenko N, Gittens R, Dvorak J. Biomechanical investigation of head impacts in football. *Br J Sports Med* 2005 Aug;39 Suppl 1:i49–i57. [PubMed: 16046356]
78. Buller DB, Andersen PA, Walkosz BJ, et al. The prevalence and predictors of helmet use by skiers and snowboarders at ski areas in western North America in 2001. *J Trauma* 2003 Nov;55(5):939–945. [PubMed: 14608169]
79. Levy AS, Hawkes AP, Hemminger LM, Knight S. An analysis of head injuries among skiers and snowboarders. *J Trauma* 2002 Oct;53(4):695–704. [PubMed: 12394869]
80. Mueller BA, Cummings P, Rivara FP, Brooks MA, Terasaki RD. Injuries of the head, face, and neck in relation to ski helmet use. *Epidemiology* 2008 Mar;19(2):270–276. [PubMed: 18277163]
81. Hagel BE, Pless IB, Goulet C, Platt RW, Robitaille Y. Effectiveness of helmets in skiers and snowboarders: case-control and case crossover study. *BMJ* 2005 Feb 5;330(7486):281. [PubMed: 15632094]
82. Cusimano MD, Kwok J. The effectiveness of helmet wear in skiers and snowboarders: a systematic review. *Br J Sports Med*. 2010 May 29;
83. Sulheim S, Holme I, Ekeland A, Bahr R. Helmet use and risk of head injuries in alpine skiers and snowboarders. *JAMA* 2006 Feb 22;295(8):919–924. [PubMed: 16493105]
84. Heintz W. The case for mandatory mouth protectors. *Phys Sportsmed* 1975 April;:61–63.
85. Winters JE Sr. Commentary: Role of Properly Fitted Mouthguards in Prevention of Sport-Related Concussion. *J Athl Train* 2001 Sep;36(3):339–341. [PubMed: 12937507]
86. Singh GD, Maher GJ, Padilla RR. Customized mandibular orthotics in the prevention of concussion/mild traumatic brain injury in football players: a preliminary study. *Dent Traumatol* 2009 Oct;25(5):515–521. [PubMed: 19614931]
87. Wisniewski JF, Guskiewicz K, Trope M, Sigurdsson A. Incidence of cerebral concussions associated with type of mouthguard used in college football. *Dent Traumatol* 2004 Jun;20(3):143–149. [PubMed: 15144445]
88. Barbic D, Pater J, Brison RJ. Comparison of mouth guard designs and concussion prevention in contact sports: a multicenter randomized controlled trial. *Clin J Sport Med* 2005 Sep;15(5):294–298. [PubMed: 16162986]
89. Labella CR, Smith BW, Sigurdsson A. Effect of mouthguards on dental injuries and concussions in college basketball. *Med Sci Sports Exerc* 2002 Jan;34(1):41–44. [PubMed: 11782645]

90. Benson BW, Meeuwisse WH. Ice hockey injuries. *Med Sport Sci* 2005;49:86–119. [PubMed: 16247263]
91. Benson BW, Hamilton GM, Meeuwisse WH, McCrory P, Dvorak J. Is protective equipment useful in preventing concussion? A systematic review of the literature. *Br J Sports Med* 2009 May;43 Suppl 1:i56–i67. [PubMed: 19433427]
92. Stevens ST, Lassonde M, de Beaumont L, Keenan JP. The effect of visors on head and facial injury in National Hockey League players. *J Sci Med Sport* 2006 Jun;9(3):238–242. [PubMed: 16716662]
93. Benson BW, Rose MS, Meeuwisse WH. The impact of face shield use on concussions in ice hockey: a multivariate analysis. *Br J Sports Med* 2002 Feb;36(1):27–32. [PubMed: 11867489]
94. Stuart MJ, Smith AM, Malo-Ortiguera SA, Fischer TL, Larson DR. A comparison of facial protection and the incidence of head, neck, and facial injuries in Junior A hockey players. A function of individual playing time. *Am J Sports Med* 2002 Jan–Feb;30(1):39–44. [PubMed: 11798994]
95. Asplund C, Bettcher S, Borchers J. Facial protection and head injuries in ice hockey: a systematic review. *Br J Sports Med* 2009 Dec;43(13):993–999. [PubMed: 19703823]
96. Benson BW, Mohtadi NG, Rose MS, Meeuwisse WH. Head and neck injuries among ice hockey players wearing full face shields vs half face shields. *JAMA* 1999 Dec 22–29;282(24):2328–2332. [PubMed: 10612320]
97. Danis RP, Hu K, Bell M. Acceptability of baseball face guards and reduction of oculofacial injury in receptive youth league players. *Inj Prev* 2000 Sep;6(3):232–234. [PubMed: 11003192]
98. Marshall SW, Mueller FO, Kirby DP, Yang J. Evaluation of safety balls and faceguards for prevention of injuries in youth baseball. *JAMA* 2003 Feb 5;289(5):568–574. [PubMed: 12578489]
99. Crisco JJ, Hendee SP, Greenwald RM. The influence of baseball modulus and mass on head and chest impacts: a theoretical study. *Med Sci Sports Exerc* 1997 Jan;29(1):26–36. [PubMed: 9000153]
100. Viano DC, McCleary JD, Andrzejak DV, Janda DH. Analysis and comparison of head impacts using baseballs of various hardness and a Hybrid III dummy. *Clin J Sport Med* 1993;4:217–228.
101. Janda DH. The prevention of baseball and softball injuries. *Clin Orthop Relat Res* 2003 Apr;409:20–28. [PubMed: 12671481]
102. Norton K, Schwerdt S, Lange K. Evidence for the aetiology of injuries in Australian football. *Br J Sports Med* 2001 Dec;35(6):418–423. [PubMed: 11726478]
103. Naunheim R, McGurren M, Standeven J, Fucetola R, Laurysen C, Deibert E. Does the use of artificial turf contribute to head injuries? *J Trauma* 2002 Oct;53(4):691–694. [PubMed: 12394868]
104. Durkin MS, Laraque D, Lubman I, Barlow B. Epidemiology and prevention of traffic injuries to urban children and adolescents. *Pediatrics* 1999 Jun;103(6):e74. [PubMed: 10353971]
105. Berg P, Westerling R. A decrease in both mild and severe bicycle-related head injuries in helmet wearing ages--trend analyses in Sweden. *Health Promot Int* 2007 Sep;22(3):191–197. [PubMed: 17728326]
106. Abu-Zidan FM, Nagelkerke N, Rao S. Factors affecting severity of bicycle-related injuries: the role of helmets in preventing head injuries. *Emerg Med Australas* 2007 Aug;19(4):366–371. [PubMed: 17655641]
107. Hagel B, Meeuwisse W. Risk compensation: a "side effect" of sport injury prevention? *Clin J Sport Med* 2004 Jul;14(4):193–196. [PubMed: 15273524]
108. Finch CF, McIntosh AS, McCrory P, Zazryn T. A pilot study of the attitudes of Australian Rules footballers towards protective headgear. *J Sci Med Sport* 2003 Dec;6(4):505–511. [PubMed: 14723399]
109. Mihalik JP, McCaffrey MA, Rivera EM, et al. Effectiveness of mouthguards in reducing neurocognitive deficits following sports-related cerebral concussion. *Dent Traumatol* 2007 Feb;23(1):14–20. [PubMed: 17227375]

Table 1
Overview of material and construction of newer football helmets as compared to an older helmet design.

| | Riddell | | | Schutt | | | Adams | | Xenith |
|----------------|--|--|--|--------------------------------|-----------------------------|---|--|--|--------|
| | Original VSR-4 | New VSR-4 | Revolution | Air Varsity Commander | DNA | Pro Elite | XI | | |
| Forehead Pad | Molded urethane | Molded urethane | Molded urethane | Dual density VN | Skydex + CF | VN | All pads utilize unique "Aware Flow" shock absorbers in a "shock bonnet" | | |
| Crown Pad | VN + CF | VN | VN + CF | Co-Molded EVA and PE | Skydex + CF | VN/EPP + EVA CF | | | |
| Back and Sides | VN + CF | VN | VN + CF | Co-Molded EVA and PE | Skydex + CF | VN/EPP + EVA CF | | | |
| Jaw Pads | Interchangeable thickness CF | Interchangeable thickness CF | VN + CF | VN + CF | VN + CF | Interchangeable thickness CF | | | |
| Fit Adjustment | Crown, back and sides padding encased in inflatable vinyl bladders | Crown, back and sides padding encased in inflatable vinyl bladders | Crown, back and sides padding encased in inflatable vinyl bladders | Halo-style tubular air bladder | Vinyl air bladder around CF | Vinyl air bladder covering crown back and sides | "Fit seeker cable" tightens the shock bonnet after chin straps are pulled tight. No pumps. | | |

Abbreviations: VN, vinyl nitrile; EPP, expanded polypropylene; CF, comfort foam; PE, polyethylene; EVA, ethylene vinyl acetate.

Adapted from Viano DC, Pellman EJ, Withnall C, Shewchenko N. Concussion in professional football: performance of newer helmets in reconstructed game impacts--Part 13 Neurosurgery. 2006 Sep;59(3):591-606